

Urge Incontinence: Diagnosis and Treatment

Urge Incontinence Symptoms

Urge incontinence is associated with a strong and overpowering urge to urinate that cannot be delayed, followed by loss of a large amount of urine. The bladder contracts without warning and urine loss occurs involuntarily. Urge incontinence is also known as overactive, spastic, or unstable bladder. If you experience urge incontinence, "When you've got to go, you've got to go!"

There is no holding time that allows you to get to the bathroom quickly enough to prevent urine leakage. Urge incontinence is the most common type of incontinence in older people.

Causes

Urge incontinence occurs when the reflexes that control the bladder are disrupted. Bladder filling signals that are transmitted to the brain are not inhibited properly in urge incontinence. This causes strong urge sensations even when the bladder is not full. The bladder contracts powerfully to the signals from the brain and the bladder outlet muscles that keep urine in the bladder cannot remain closed. Involuntary urine leakage results.

While a number of health problems or other conditions may be associated with urge incontinence, in many instances the direct cause is unknown. In some cases, urge incontinence may be the result of conditions such as:

- Neurologic problems — Diabetes Mellitus, Parkinson's disease, stroke, spinal chord injuries, multiple sclerosis, dementia and Alzheimer's disease
- Urinary tract infection
- Urethritis or vaginitis
- Kidney or bladder stones or tumors
- Interstitial cystitis
- Enlarged prostate, tumor, or prostatitis
- Decreased estrogen in the post-menopausal woman
- Previous bladder or pelvic surgery
- Side effect from diuretics (water pills)

Other conditions that may lead to urge incontinence include...

- Constipation
- Bladder irritants such as caffeine
- Dehydration

(Adapted with permission from the Practicing Physician Education Project 2006)

Urge Incontinence: Diagnosis and Treatment

Causes

(Continued)

Diagnosis

- Excess urine production

The primary goal of diagnosis is to identify the causes and the factors contributing to urge incontinence. Often there are multiple causes. Initial evaluation, therefore, includes the following components:

History. Your health care provider needs to find out about the circumstances associated with the onset of urge incontinence and to identify the presence of any health conditions that affect urinary tract function.

Bladder Diary. This is used to determine the timing, symptoms, and associated or precipitating factors of the incontinence.

Physical Exam. An exam of your major body systems is performed to check for potential causes of incontinence.

Urinalysis and Urine Culture and Sensitivity. These tests are performed to evaluate urinary system functioning and to test for urinary tract infection.

Simple Cystometry. This is used to assess the function of the bladder when it is filled with fluid.

Stress Maneuver. This is used to screen for incontinence during a cough or sneeze with a full bladder.

Other laboratory and radiologic testing may be required to assist in diagnosis.

Treatment

There are a number of treatments that may be used alone or together to treat urge incontinence. They include:

- Bladder training
- Identification and elimination of bladder irritants
- Bowel management to prevent constipation and fecal impaction
- Pelvic muscle exercises
- Bladder urge control
- Medication: Anticholinergic, antidepressants, and hormone replacement therapy

Treatment is tailored to the needs, symptoms, capabilities and preferences of the individual.

(Adapted with permission from the Practicing Physician Education Project 2006)